Determinants of Domestic Water Consumption: A Case Study

in Northern Portugal

<u>A. Manuela Gonçalves</u>^{1*}, Cristina Matos ², Ana Briga-Sá ³, Sandra Pereira ⁴, Isabel Bentes ⁵ and Diana Faria⁶

¹CMAT – Centro de Matemática, DMA-Departamento de Matemática e Aplicações, Universidade do Minho, Portugal ^{2,3,4,5,6}C-MADE – Centro de Materiais e Tecnologias Construtivas, Universidade da Beira Interior, ECT – Escola de Ciências e Tecnologia, Universidade de Trás-os-Montes e Alto Douro, Portugal

> <u>mneves@math.uminho.pt</u> (*corresponding and presenting author) <u>crismatos@utad.pt</u>

Poster Abstract

Efficient management of water resources, in both rural and urban areas, requires a full understanding of existing patterns of water use. Water demand management has been mainly focused on meeting agriculture water demand, whereas domestic water demand is largely ignored, and household water consumption has not been thoroughly researched in the majority of the countries. The World Health Organization (WHO) defined "domestic water" as water used for all domestic purposes including consumption, bathing and food preparation [1], [5]. Information regarding domestic water consumption is vital but is still lacking. The success of domestic water demand management strategies depends on identifying the determinants, and their interaction, that influence water consumption at a household scale [3].

This paper presents an empirical analysis of domestic water consumption and factors influencing water consumption in Vila Real County, in Northern Portugal. Through a field survey, the data were collected from December 2016 to January 2017 from 245 urban and rural households in 20 parishes of Vila Real County, and determinants influencing domestic water consumption are studied. Data analysis was performed by descriptive statistics, non-parametric tests and ordinal regression, namely by comparing the two groups (urban and rural households) [2], [4].

Keywords: Survey, Domestic Water Consumption, Non-parametric Tests Analysis, Ordinal Regression.

Acknowledgements

This work was partially supported by the Research Centre of Mathematics of the University of Minho with the Portuguese Funds from the "FCT - Fundação para a Ciência e a Tecnologia", through the Project PEstOE/MAT/ UI0013/2017. This work was partially funded by POCI-0-0145-FEDER-016730 Project (PTDC/AAG-REC/4700/2014) with the designation ENERWAT: Water to energy: characterization, modelling and measures for the reduction of urban and rural household consumption, financed by the Foundation for Science and Technology and co-financed by the European Regional Development Fund (FEDER) through the COMPETE 2020 - Programme Operational Competitiveness and Internationalization (POCI). This work was partially supported by the FCT (Portuguese Foundation for Science and Technology) through the project PEst-OE/ECI/UI4082/2013 (C-MADE).

References

- [1] World Heatlth Organization (2002). *Guidelines for Drinking-Water Quality: Addendum Microbiological Agents in Drinking Water*, 2nd ed.; WHO: Geneva, Switzerland.
- [2] Higgins J.L. (2004). Introduction to Modern Nonparametric Statistics. Toronto, Canada: Thomson.
- [3] Loureiro D., Pinheiro L., Rebelo M., Salgueiro A.R., Medeiros N., Covas D., Alegre H. (2008). Estudo dos fatores mais relevantes que influenciam o consumo doméstico de água: O caso de estudo do complexo de edifícios: Twin-Towers. LNEC.
- [4] O'Connel A.A. (2006). *Logistic Regression Models for Ordinal Response Variables*. Series: Quantitative Applications in the Social Sciences. SAGE Publications, Inc., London.
- [5] Vieira A.S., Ghisi E. (2016). Water-energy nexus in houses in Brazil: comparing rainwater and gray water use with a centralized system. *Water Science and Technology: Water Supply* 16 (2), 274-283.
- © ESTG, Instituto Politécnico do Porto